



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME

New names and the final members of new combinations are in **heavier type**.

- Abies grandis*, 99; *Mertensiana*, 100; *Pattoniana*, 100
Acalypha jamaicensis, 7; *pruinosa*, 7
Acer platanoides, 408; *rubrum*, 407; *saccharinum*, 408
Aconiopteris gorgonea, 572; *obtusata*, 572
Acorus Calamus, 107
Acrophorus Macraeanus, 582; *repens*, 582
Acrostichum, 258, 571; *aemulum*, 571, 572; *calomelanos*, 588; *crassifolium*, 572; *gorgonium*, 572; *Helleri*, 571; *hirtum*, 570; *lineatum*, 275; *micans*, 570; *micradenium*, 570; *pellucidomarginatum*, 572; *reticulatum*, 572; *Spicant*, 275; *squamosum*, 570; *vestitum*, 570; *Wawrae*, 571
Actinostemon jamaicensis, 7
Adiantum, 568, 577; *Bennettii*, 577; *Capillus*, 577; *Capillus-veneris*, 577; *chinense*, 585; *chusanum*, 585; *emarginatum*, 577; *Levingei*, 577; *tenuifolium*, 585; *Wattii*, 577.
Adnaria, 550
Aenothera grandiflora, 519, 521; *longiflora*, 521
Ailanthus glandulosa, 420
ALDEN, I. A contribution to the life history of *Uvularia sessilifolia*, 439
Alectoria jubata, 201; *pubescens*, 201
Alisma brevipes, 101; *Geyeri*, 101; *Plantago-aquatica*, 101
Alismaceae, 101
Alliaceae, 110
Allium, 16, 19, 22; *Cepa*, 15, 17, 20
Allosorus aquilinus, 575; *ternifolius*, 576
Alnus, 439
Alopecurus aristulatus, 102; *fulvus*, 102; *pallescens*, 102
Alsinaceae, 314
Alsine alpestris, 315; *Edwardsii*, 315; **Palmeri**, 315; *stricta*, 315; *strictiflora*, 315; *subvestita*, 315
Alsinopsis dawsonensis, 316; **pusilla**, 316
Amaranthaceae, 313
Amaranthus californicus, 313; *carneus*, 313; *gracilis* *pubescens*, 313; **pubescens**, 313
Ammodenia oblongifolia, 317
Ampelocissus Alexandri, 12
Ampelopsis, 39, 50
Anacardiaceae, 423
ANDREWS, F. M. Protoplasmic streaming in *Mucor*, 455
Andromeda Novae-caesareae, 390, 405; *Pfaffiana*, 388; *Snowii*, 390, 405
Androvettia, 389
Anetium reticulatum, 572
Anticlea, 108; *coloradensis*, 108; *elegans*, 108; *porrifolia*, 108, 109; **vaginata**, 108, 109
Apios Apios, 80
Aphanolejeunea, 222, 224, 329
Arabis brevisiliqua, 326; *lignifera*, 326; **MacDougallii**, 326; *subpinnatifida*, 326
Araceae, 107
Aralia cottondalensis, 403; *decurrens*, 404; *Saportana*, 403, 404; *Wellingtoniana*, 390, 402-404; *Wellingtoniana* *Vaughanii*, 388
Araucaria bladenensis, 389; *Jeffreyi*, 389
Arenaria cephaloidea, 316; *congesta*, 316; *dawsonensis*, 316; *diffusa*, 316; *Fendleri* *diffusa*, 316; *laricifolia*, 316; **laxiflora**, 316; **lithophila**, 316; *peplodes major*, 317; *pusilla*, 316; *sitchensis*, 317; *subcongesta* *lithophila*, 316
Aristida bromoides, 102; *fasciculata* *Hookeri*, 102; *oligantha*, 102; *longiseta*, 102
Arrhenatherum elatius, 103
Artabotrys odoratissima, 502
Arthrotaxopsis, 393
Ascobolaceae, Methods of culture and the morphology of the archicarp in certain species of the, 139
Ascobolus, 140, 141, 144-146, 149, 150, 167, 170, 171, 187, 188; *aerugineus*, 151; *albidus*, 144, 148, 149; *amoenus*, 179; *atrofuscus*, 171; *brunneus*, 179; *carbonarius*, 141, 148, 149, 151, 159-163, 170-172, 187, 189-195; *carbonicola*, 171; *denudatus*, 143, 149; *furfuraceus*, 141, 144-149, 151, 168, 169, 185, 186, 192-194; *glaber*, 146, 151, 168, 169, 180, 193, 194; *immersus*, 142, 149, 151, 164, 165, 169, 184, 186, 190, 193, 194; *Leveillei*, 151, 166, 169, 179, 180, 194; *Leveillei americanus*, 152, 167, 169, 179; *mirabilis*, 191; *perplexans*, 144, 148; *pulcherrimus*, 457; *pusillus*, 151, 152, 170, 172, 193; *sp.*, 148, 151, 156; *stictoides*, 179; *viridis*, 141, 148, 151, 152, 164, 171, 172, 193; *viridulus*, 152, 170; *Winteri*, 151, 152, 154, 156, 158, 179, 180, 181, 184, 186, 190, 193, 195; *xylophilus*, 167, 169, 193

- Ascodesmis*, 139, 186, 191; *nigricans*, 141, 145, 148, 152, 169, 170
Ascophanus, 187; *Aurora*, 152; *carneus*, 144, 147-149, 151, 152, 169, 178-180, 186, 190, 193, 457, 491; *glaucellus*, 152; *granuliformis*, 152; *Holmskjoldii*, 152; *lacteus*, 152; *microsporus*, 152; *minutissimus*, 152; *ochraceus*, 146, 152, 190; *pilosus*, 142; *saccharinus*, 143, 148; *sarcobius*, 152, 169, 178, 193
Asparagin, effect of, on absorption and growth in wheat, 429
Aspergillus, 149, 190; *niger*, 535
Aspidium cordifolium, 583; *cyatheoides*, 599; *Ecklonii*, 598; *exaltatum*, 584; *Filix-mas fusco-atrum*, 592; *glabrum*, 592; *glabrum pusillum*, 593; *glabrum quadripinnatum*, 592; *globuliferum*, 591; *hawaiiense*, 594; *Hudsonianum*, 601; *juglandifolium macrosorum*, 26; *latifrons*, 593; *molle*, 599; *nymphale*, 599; *paleaceum*, 591; *parallelogrammum*, 591; *parasiticum*, 599; *patens*, 599; *propinquum*, 598; *resiniferum*, 598; *rubiginosum*, 594; *Schkuhrii*, 584; *Speluncae*, 586; *squamigerum*, 594; *tuberosum*, 583; *undulatum*, 583; *unitum*, 598; *violascens*, 599
Asplenium, 258, 569; *aquilinum*, 575; *Spicant*, 275
Athyrium, 569
Atriplex aptera, 311; ***buxifolia***, 311; *canescens*, 311, 312; *confertifolia*, 312; ***Garretti***, 312; *Nuttallii*, 311; *occidentalis*, 312; *odontophora*, 311; *phyllostegia*, 312; ***tetraptera***, 311, 312
Avena americana, 103; *Hookeriana*, 103; *Mortoniana*, 103
Azolla, 232

Baptisia tinctoria, 69
Batodendron, 549, 553
BEATTIE, J. H., and SKINNER, J. J.
Effect of asparagin on absorption and growth in wheat, 429
Benzoin venustum, 388, 390, 399
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII. Texas, 387; Notes on the genus *Widdringtonites*, 341
BICKNELL, E. P. The ferns and flowering plants of Nantucket—IX, 69; —X, 415
Bicornia, 553
Bidens clarendonensis, 9
Blechnum, 257-259, 261, 276; *attenuatum*, 272; *boreale*, 275; *capense*, 367, 375; *Christii*, 362; *costaricensis*, 363; *danaeaceum*, 364; *dodioides*, 275, 277; *ensiforme*, 263; *falciforme*, 365; *Giesbreghtii*, 277; *L'Herminieri*, 268; *lineatum*, 367; *onocleoides*, 271; *ornifolium*, 371; ***Plumieri***, 269; *polypodioides*, 272; *sessilifoli[um]*, 373; *Spicant*, 275; *stoloniferum*, 277; *striatum*, 375; *tubulare*, 377; *varians*, 379; *violaceum*, 379; *Werckleanum*, 382
Blepharidachne Kingii, 103
Boryana [*Struthiopteris*], 378
Botrychium, 234; *subbifoliatum*, 234
Boudiera, 152; *Claussenii*, 152, 170
Brachyphyllum crassicaule, 393; *macrocarpum formosum*, 389, 390, 392, 406; *obesiforme elongatum*, 393
BRAINERD, E. Violet hybrids between species of the *palmata* group, 85
Brassicaceae, 322
BRITTON, N. L. Studies of West Indian plants—IV, 1
Briza maxima, 104
BROADHURST, J. The genus *Struthiopteris* and its representatives in North America, 257, 357
Brodiaea, 110, 111; *capitata*, 111; *capitata pauciflora*, 111; *congesta*, 111; *grandiflora*, 110, 111; *insularis*, 111; *volubilis*, 111

Caeoma nitens, 113-116
Caesalpiniaceae, 69
Calamagrostis, 103; *lucida*, 103; *rubescens*, 103; *Vaseyi*, 103
Callitrichaceae, 422
Callitriche heterophylla, 422
Callitris, 342, 343
Calyptridium roseum, 314
Canangium Brandesianum, 504; *odoratum*, 504
Cameraria angustifolia, 5; *latifolia*, 5, 6; ***microphylla***, 6; ***oblongifolia***, 6; *retusa*, 5
Caryophyllaceae, 317
Case of changed polarity in *Spirogyra elongata*, A, 509
Cassia Chamaechrista, 69
Cathartolinum intercursum, 418
Celtis Douglasii, 305; *occidentalis*, 305; *reticulata*, 304, 304; ***rugosa***, 304
Cerastium arvense fuegianum, 314, 315; *fuegianum*, 314; *strictum*, 315; ***thermale***, 314
Ceropteris, 568, 588; *calomelanos*, 588; ***ochracea***, 588
Cetraria nivalis, 201; *islandica*, 201
Chara, 467
Cheilanthes, 579; *Lidgatii*, 579; *ternifolia*, 576
Cheilolejeunea decidua, 279
Cheiranthus, 323; *argillosus*, 324; *aridus*, 323, 324; *asper*, 323; *asperimus*, 324; *Bakeri*, 324; *cheiranthoides*, 323; *elatus*, 323; *inconspicuus*, 323; *nivalis*, 324; *nivalis amoenus*, 324; *Pallasii*, 324; *pygmaeus*, 324; *syrticola*, 323; *Wheeleri*, 324
Cheirinia, 323; ***amoena***, 324; ***argillosa***,

- 324; *arida*, 323; *aspera*, 323, 325; *asperrima*, 324; *Bakeri*, 324; *brachycarpa*, 325; *cheiranthoides*, 323; *elata*, 323; *inconspicua*, 323; *nivalis*, 324; *oblanceolata*, 324, 325; *Pallasii*, 324; *radicata*, 324; *syrticola*, 323; *Wheeleri*, 324
 Chenopodiaceae, 310
 Chenopodium calceoliforme, 313; desiccatum, 311; lanceolatum, 311; leptophyllum, 310; oblongifolium, 310; opulifolium, 311; paganum, 311; *pratericola*, 310; rubrum, 311; succosum, 311; viride, 311
 Chiogenes, 550
 Chloris brevispica, 103
 Chorizanthe brevicornu, 310; *spathulata*, 309
 Cibotium, 242; Chamissoi, 242, 243; glaucum, 243, 244; Menziesii, 242, 243; pruinatum, 243
 Cinna arundinacea, 103
 Cinnamomum, 400; ellipsoideum, 388; Heerii, 388; membranaceum, 390, 401; sp., 388
 Cincinialis aquilina, 575
 Cissus, 39, 41, 50
 Cladophora, 509
 Clerodendron calcicola, 9; spinosum, 10
 Clusia clarendonensis, 7
 Cocculus, 400
 Codorolla, 554
 Cololejeunea, 210-212, 222-234, 329; Biddlecomiae, 223; floccosa, 607; peraffinis, 608; platyneura, 608
 Colura, 209, 210
 Colurolejeunea, 329
 Colutea primordialis, 390, 396
 Commelinaceae, 107
 Coniogramme, 568, 569; fraxinea, 589
 Contribution to the life history of Uvularia sessilifolia, A, 439
 Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII. Texas, 387
 Coprinus, 186
 Corema Conradii, 422
 Cornophyllum vetustum, 390, 404
 Coronaria, 317
 Coronilla varia, 74
 Corylus, 439
 Cracca virginiana, 74
 Cuba, Undescribed species from, 9
 Cuba, The genus Ginoria in, 12
 Cubonia sp., 152
 Cunninghamites, 389
 Cyatheaceae, 237, 241
 Cyclolejeunea, 329
 Cymbopetalum penduliflorum, 502
 Cyrtomium, 569
 Cystopteris Douglasii, 587; sandwichensis, 587
 Cystopus, 188
 Cytisus scoparius, 70
 DACHNOWSKI, A. The relation of Ohio bog vegetation to the chemical nature of peat soils, 53
 Danthonia spicata, 103
 DARLING, C. A. Mitosis in living cells, 407
 Davallia biflora, 584; chinensis, 585; chusana, 585; flaccida, 586; Khasiyana, 586; Macraeana, 582; Mannii, 582; polypodioides, 585; remota, 585; Spelunca, 586; strigosa, 586; tenuifolia, 585
 DAVIS, B. M. Was Lamarck's evening primrose (Oenothera Lamarckiana Seringe) a form of Oenothera grandiflora Solander?, 519
 Delphinium bicolor, 320, 322; *coelestinum*, 320; Geyeri, 322; *Helleri*, 321; *Leonardi*, 320; multiflorum, 322; reticulatum, 322; scaposum, 321; scopulorum, 322; *viscidum*, 322; *xyloirrhizum*, 321
 Dendropanax arboreum, 1, 2; *blakeanum*, 4; *brachypodium*, 2; *cordifolium*, 4; *cuneifolium*, 2; *elongatum*, 3; *grande*, 4; *grandiflorum*, 3; *laurifolium*, 2; *nutans*, 2; *pendulum*, 2, 4; *samydifolium*, 1
 Dendropanax in the West Indies, 1
 Dennstaedtia strigosa, 586
 Deschampsia caespitosa, 103; *pungens*, 103
 Desmos, 504, 505; chinensis, 502, 504; cochinchinensis, 504-506; *crinitus*, 507; *dasyaschalus*, 507; *desmanthus*, 508; *dumosus*, 506; *Dunalii*, 506; *elegans*, 506; *Lawii*, 506; *longiflorus*, 507; *pannosus*, 506; *praecox*, 507; *stenopetalus*, 507; *viridiflorus*, 506; *Wrayi*, 507; *zeylanicus*, 506
 Desmos the proper generic name for the so-called Unonas of the Old World, 501
 Detonia trachycarpa, 170
 Diastoloba, 280
 Dichasium parallelogrammum, 591
 Dichelostemma, 110, 111; congesta, 111; multiflora, 111
 Dicksonia glauca, 244; japonica, 585; Menziesii, 243; splendens, 243; strigosa, 585
 Dicranopteris, 238, 239; dichotoma, 241; *emarginata*, 239, 240; glabra, 239, 240; glauca, 239, 240; linearis, 239, 240; *owhyhensis*, 239, 241
 Didymoglossum humile, 246
 Diellia, 568, 580; Alexandri, 580, 581; *centifolia*, 580; erecta, 580, 581; falcata, 580, 581; Knudsenii, 580, 582; *laciniata*, 580, 581; *Mannii*, 580, 582; *pumila*, 580
 Diospyros primaeva, 388; Steenstrupi, 388

- Diplasiolejeunea, 209, 210, 212, 222, 224, 329, 603, 605; *armatiloba*, 216; *brachyclada*, 216, 217, 220, 222, 225, 605; *Johnsonii*, 603-606, 611; *pellucida*, 210-225, 605, 606; *pellucida malleiformis*, 215, 222; *Rudolphiana*, 220-222, 224, 225, 605; *unidentata*, 217, 219, 221, 222, 224, 225, 605
- Diplasiolejeunea, Vegetative reproduction in, 222
- Diplazium fraxineum, 589
- Diplusodon ginorioides, 13
- Dipterostemon, 110; *capitatus*, 111; *insularis*, 111; *pauciflorus*, 111; *pulchellus*, 111
- Discoid gemmae in *Radula*, 329
- Disterigma, 552
- DODGE, B. O. Methods of culture and the morphology of the archicarp in certain species of the Ascobolaceae, 139
- Dondia calceoliformis*, 313
- Doodia, 276, 569
- DORSEY, M. J. Variation in the floral structures of *Vitis*, 37
- Doryopteris, 567, 576; *decipiens*, 576; *decora*, 576, 577; *pedata*, 576
- Draba andina*, 327, 328; *aurea*, 328; *aureiformis*, 328; *decumbens*, 328; *deflexa*, 328; *densiflora*, 327, 328; *glacialis pectinata*, 327; *lapilutea*, 328; *luteola*, 328; *pectinata*, 327, 328; *praealta*, 328; *reflexa*, 328; *surculifera*, 328; *uber*, 328; *yellowstonensis*, 328
- Dryopteris, 568, 569, 589; *acutidens*, 590, 597; *crinalis*, 590, 595; *cyatheoides*, 591, 599-601; *dilatata*, 597; *fijiensis*, 595; *Filix-mas*, 592, 594; *fusco-atra*, 589, 592; *glabra*, 590, 592, 594; *globulifera*, 589, 591, 599; *hawaiiensis*, 590, 594; *honolulensis*, 590, 595; *Keraudreniana*, 590, 596; *latifrons*, 590, 593, 595; *nuda*, 590, 592, 593; *paleacea*, 589, 591, 592; *parasitica*, 591, 598, 599; *parvula*, 590, 593; *propinqua*, 590, 598; *pusilla*, 593; *rubiformis*, 590, 596; *rubiginosa*, 590, 594; *sandwicensis*, 590, 598, 600; *spinulosa*, 592, 597; *squamigera*, 590, 594; *stegnogrammoides*, 591, 600, 601; *truncata*, 591, 601; *unidentata*, 590, 597
- Effect of asparagin on absorption and growth in wheat, 429
- Effect of guanidin on plants, 535
- Eichhornia*, 16, 19, 21, 22; *speciosa*, 15
- Elaeagia cubensis*, 10
- Elaphoglossum*, 567, 569, 571; *aemulum*, 570, 571; *conforme*, 569; *gorgonium*, 570, 572; *hirtum*, 570; *micradenium*, 570; *nitidum*, 570; *squamosum*, 570; *reticulatum*, 570, 572; *vestitum*, 570; *Wawrae*, 570, 571
- Elisanthe*, 318; *Drummondii*, 318
- Elymus*, 107
- Empetraceae, 422
- Endolepis phyllostegia*, 312
- Ephedra nevadensis*, 100; *viridis*, 100
- Ephedraceae, 100
- Eragrostis hypnoides*, 104; *lutescens*, 104; *neomexicana*, 104; *secundiflora*, 104
- Eremochloa Kingii*, 103
- Ericaceae, 553
- Ericales, 405
- Eriocoma cuspidata*, 102; *hymenoides*, 102
- Eriogonum*, 308, 309; *androsaceum*, 309; *angulosum*, 309; *aureum*, 308, 309; *Baileyi*, 309; *bicolor*, 309; *biumbellatum*, 306; *campanulatum*, 307; *chloranthum*, 308; *clavellatum*, 309; *compositum*, 309; *corymbosum*, 308; *crassifolium*, 308; *croceum*, 307; *deflexum*, 309; *densum*, 309; *depauperatum*, 309; *depressum*, 308; *dichotomum*, 309; *elatum*, 309; *flavum*, 308; *Hookeri*, 309; *idahoense*, 307; *insigne*, 309; *leptophyllum*, 309; *longilobum*, 309; *Mearnsii*, 309; *micranthum*, 309; *microthecum*, 307; *neglectum*, 307; *nidularium*, 309; *nutans*, 309; *ochrocephalum*, 309; *Ordii*, 309; *orendense*, 307; *ovalifolium depressum*, 308; *Parryi*, 309; *polifolium*, 309; *Porteri*, 309; *proliferum*, 309; *puberulum*, 309; *pulvinatum*, 309; *racemosum*, 309; *ramosissimum*, 308, 309; *rubidum frigidum*, 308; *salicinum*, 308; *Shockleyi*, 309; *spathulatum*, 308, 309; *spathuliforme*, 307; *spergulinum*, 309; *strictum*, 308, 309; *subreniforme*, 309; *sulcatum*, 309; *tenellum*, 309; *Thomasii*, 309; *Thompsonae*, 309; *thymoides*, 309; *trichopodium*, 309; *tristichum*, 308; *turbidatum*, 309; *umbellatum*, 307; *umbelliferum*, 307; *villiflorum*, 309; *vimineum*, 309; *Wetherillii*, 309; *Wrightii*, 309
- Erodium cicutarium*, 417
- Erysimum*, 323; *alpestre*, 324; *amoenum*, 324; *argillosum*, 324; *asperum*, 323; *asperum alpestre*, 324; *asperum inconspicuum*, 323; *Bakeri*, 324; *cheiranthoides*, 323; *elatum*, 323; *inconspicuum*, 323; *nivale*, 324; *oblanceolatum*, 324; *officinale*, 323; *parviflorum*, 323; *pygmaeum*, 325; *radicatum*, 324; *syrticum*, 323; *Wheeleri*, 324
- Erythronium*, 441
- ESTABROOK, A. H., and LIVINGSTON, B. E. Observations on the degree of stomatal movement in certain plants, 15
- Eudianthe*, 317
- Eucallitris*, 342
- Eucalyptus Geinitzi*, 390, 402
- Eugenia primaeva*, 388
- Euphorbia Cyparissias*, 422; *maculata*, 421; *polygonifolia*, 421; *Preslii*, 421

- Euphorbiaceae, 421
 Eupteris aquilina, 575
 Eurotia lanata, 312; **subspinosa**, 312, 313
 EVANS, A. W. Hepaticae of Puerto Rico—XI. Displasiolejeunea, 209; New West Indian Lejeuneae—II, 603
- Falcata monoica, 80; Pitcheri, 80
 Ferns and flowering plants of Nantucket, The—IX, 69; —X, 415
 Ferns from Tropical America, New, 285
 Festuca, 105, 106; campestris, 105; confinis, 106; dasyclada, 105; Kingii, 106; megalura, 105; ovina calligera, 105; pacifica, 105; reflexa, 105; scabrella, 105; Thurberi, 105; viridula, 105; Watsonii, 106
 Ficus daphnogenoides, 390, 394; glaucocoeana, 388
 Filix, 568, 587; **Douglasii**, 587
 Filicales, 232, 236
 Flax rust, 113
 Frenela, 342
 Frenelites, 343
 Frenelopsis, 343, 346, 347; bohémica, 346; Hoheneggeri, 346; ramosissima, 346, 347
 FROMME, F. D. Sexual fusions and spore development of the flax rust, 113
 Frullania, 279, 280, 284, 329; Asagrayana, 280; Bolanderi, 279, 284; Brittoniae, 280, 283, 284; cleistostoma, 284; dilatata, 280, 283; eboracensis, 279, 280, 281, 283, 284; fragilifolia, 280; inflata, 280; plana, 280, 283, 284; riparia, 280, 283, 284; saxicola, 280; Selwyniana, 280; squarrosa, 221; Tamarisci, 280
 Frullaniae, Vegetative reproduction in the New England, 279
 Funkia, 19, 22; ovata, 15
- Gaultheria, 550
 Gaylussacia, 549, 550; buxifolia, 550; dumosa, 551; frondosa, 555; nana, 555; tomentosa, 555
 Genus Cameraria, The, 5
 Genus Ginoria in Cuba, The, 12
 Genus Struthiopteris and its representatives in North America, The, 257, 357
 Geraniaceae, 415
 Geranium carolinianum, 415; maculatum, 415; molle, 416; pusillum, 416; Robertianum, 415
 Gilibertia brachypoda, 2
 Ginoria americana, 12, 13; **arborea**, 13; curvispina, 13; Diplusodon, 13; **ginorioides**, 13; spinosa, 13
 Gleditsia triacanthos, 69
 Gleichenia glauca, 240; hawaiiensis, 241; linearis, 241; longissima, 240; owhyhensis, 241
 Gleicheniaceae, 237, 238
 Gonocormus saxifragoides, 245
 Guanidin, the effect of, on plants, 535
 Guatteria, 502
 Gymnoascus, 186; ruber, 186
 Gymnoconia interstitialis, 113, 114, 116
 Gymnogramme calomelanos, 588; javanica, 589; pilosa, 589; tartarea, 588
 Gymnospermae, 391
 Gymnosporangium clavariaeforme, 119, 128
 Gymnotheca Douglasii, 236
- Hawaiian Islands, A taxonomic study of the Pteridophyta of the, 227, 567
 Hedera nutans, 3, 4
 Hepatica, 439
 Hepaticae of Puerto Rico—XI. Diplasiolejeunea, 209
 Hesperis pygmaeus, 325
Hesperochloa, 106; **Kingii**, 106
 Hesperopeuce **Mertensiana**, 100; Pattoniana, 100
 Hexaclinis, 342
 Hicriopteris speciosa, 240
 Homotropantha, 280
 Honckenya oblongifolia, 317
 Hookera, 111; coronaria, 110; pulchella, 111
 Hordeum Aegiceras, 106; caespitosum, 106; montanense, 106; murinum, 106
 HOWE, R. H. JR. The lichens of the Linnean Herbarium with remarks on Acharian material, 199
 Hugeria, 551
 Humaria, 190, 191; granulata, 191, 195
 Humata Mannii, 582
 Hymenodium crassifolium, 572; reticulatum, 572
 Hymenophyllum, 244, 245, 247; Baldwinii, 247; lanceolatum, 247, 248; obtusum, 247, 248; recurvum, 247; tumbidgeense, 247
 Hymenophyllaceae, 237, 244
 Hypocopra, 190
 Hypolepis, 568, 578; **flaccida**, 578, 579; punctata, 578; tenuifolia, 578, 579
- Ilex, 387; bronxensis, 426, 427; **fastigiata**, 426; glabra, 425; verticillata, 426, 427; verticillata padifolia, 426
 Illicaceae, 425
 Index to American Botanical Literature, 29, 81, 133, 205, 249, 293, 349, 411, 447, 511, 561, 613
 Inga cretacea, 388
 Isatis, 20, 22; tinctoria, 15
- Jamaica, Undescribed species from, 7
 Juncaceae, 110
 Juncus arizonicus, 110; columbianus, 110; ensifolius, 110; Jonesii, 110; mexi-

- canus, 110; Regelii, 110; saximontanus, 110; Tracyi, 110; uncialis, 110
 Jungermannia floccosa, 607; pellucida, 213, 214; unidentata, 217, 219
 Juniperaceae, 99
 Juniperites baccifera, 342
 Juniperus, 281; horizontalis, 100; Knightii, 99; monosperma, 99; prostrata, 100; Sabina procumbens, 100; utahensis, 99, 100; virginiana, 99
 Lachnea melaloma, 170; stercorea, 147, 190
 Larix decidua, 408
 Lasiodolus, 187; equinus, 142, 152, 168, 169; pulcherrimus, 140
 Lastraea glabra, 592; globulifera, 591; latifrons, 593; rubiginosa, 594; squamigera, 594; truncata, 591
 Lathyrus maritimus, 79; pilosus, 79; tuberosus, 80
 Laurophyllum minus, 390, 402
 Laurus, 387; plutonia, 390, 401; proteaefolia, 388
 Leiolejeunea grandiflora, 603
 Lejeunea, 210, 329, 603; albifolia, 214; floccosa, 607; ocellulata, 213, 214; pellucida, 213; platyneura, 608; unidentata, 217, 220
 Lemna cyclostosa, 107; minima, 107; minor, 107; perpusilla, 107
 Lemnaceae, 107
 Lepidium hirsutum, 322; intermedium pubescens, 322; medium pubescens, 322; virginicum texanum pubescens, 322
 Leptocolea, 210, 222-224, 329, 603, 607; appressa, 606-608; floccosa, 607, 608; scabriflora, 607
 Lespedeza, 75; Bicknellii, 77; Brittonii, 76; capitata, 78; frutescens, 77; hirta, 77; Nuttallii, 76; procumbens, 75; Stuvei, 76; velutina, 77; violacea, 77; virginica, 76
 Letharia vulpina, 201
 Lichens of the Linnean Herbarium with remarks on Acharian material, The, 199
 Limnia depressa, 314; spathulata, 314; utahensis, 314
 Linaceae, 418
 Linderia venusta, 399
 Lindsaya Alexandri, 581; centifolia, 580; chinensis, 585; Knudsenii, 582; laciniata, 581; laciniata subbipinnata, 581; Mannii, 582; repens, 583
 Linum, 419; floridanum, 419, 420; intercursum, 418-420; medium, 419; striatum, 418; sulcatum, 420; usitissimum, 118, 418; virginianum, 418
 Liquidambar integrifolium, 388
 Liriodendron pinnatifidum, 388, 395; quercifolium, 390, 395; Snowii, 395
 Litobrochia decora, 577; decipiens, 576
 Litsea falcifolia, 399
 LIVINGSTON, B. E., and ESTABROOK, A. H. Observations on the degree of stomatal movement in certain plants, 15
 Lomaria, 257, 259; acrodonta, 371, 373; attenuata, 272, 383; Boryana, 377; brasiliensis, 375, 377; costaricensis, 363; crenata, 275, 277; danaeacea, 364, 365; decrescens, 271, 273; deflexa, 365; divergens, 271; ensiformis, 263; exaltata, 264, 266, 271; falciformis, 365; Féei, 272; fragilis, 271, 273; Gheisbreghtii, 277; L'Herminieri, 266, 267; lineata, 366; longifolia, 371, 372; martinicensis, 269; mexicana, 271, 273; mucronata, 266, 271; onocleoides, 271; ornifolia, 373; Plumieri, 269; poly-podioides, 271; procera, 366, 383, 384; pteropus, 274; robusta, 369, 370; rufa, 369, 383; Ryani, 375, 377, 383; Schiedeana, 370-373; Schomburgkii, 383; sessilifolia, 373; spectabilis, 371, 372, 380; Spicant, 275; Spicant serratus, 277; spissa, 362; stolonifera, 277; striata, 375; tuberculata, 375; varians, 379; violacea, 379; Werckleana, 382
 Lomaridium Herminieri, 267; Plumieri, 269
 Lonchitis tenuifolia, 578
 LORENZ, A. Vegetative reproduction in the New England Frullaniae, 279
 Loxsomopsis, 285; costaricensis, 286; Lehmanii, 286; notabilis, 285, 288
 Lychnis, 317, 318; affinis, 318; apetal, 319; attenuata, 319; Drummondii, 318; Kingii, 319; montana, 318; Parryi, 319; striata, 318; Taylorae, 318; triflora, 318
 Lycopodiales, 232
 Lysichiton camtschaticense, 107
 MACKENSEN, B. Three new species of Opuntia, with a discussion of the identity of Opuntia Lindheimeri, 289
 Magnolia Boulayana, 388; speciosa, 388, 390, 395
 Malapoenna falcifolia, 390, 399
 Malvales, 399
 Marattia, 236; alata, 236; Douglasii, 236
 Marattiaceae, 236
 Marattiales, 232, 235
 Marsilea crenulata, 233; natans, 233; quadrifolia, 233; villosa, 233
 Marsileaceae, 232, 233
 Matteuccia, 259
 MAXON, W. R. Notes on the North American species of Phanerophlebia, 23
 Maytenus clarendonensis, 8
 Medicago lupulina, 72; sativa, 71
 Meibomia canadensis, 75; obtusa, 74; rigida, 75; sessilifolia, 75

- Melampsora* Lini, 113, 118, 120-122, 127, 128; *Rostrupi*, 113, 120
Melandrium, 317, 318; *affine*, 318; *apetalum*, 319; *trifolium*, 318
Melanthaceae, 108
Melilotus alba, 72; *officinalis*, 72
Melodorum, 503; *fruticosum*, 503; *latifolium*, 504
Mertensia dichotoma, 241; *emarginata*, 240; *excelsa*, 240; *glabra*, 240; *glauca*, 239; *hawaiiensis*, 241; *pinnata*, 240
Mespilodaphne, 401
Methods of culture and the morphology of the archicarp in certain of the *Ascobolaceae*, 139
Mettenia acutifolia, 9; *globosa*, 9
Metzgeria, 279, 329
Metzgeriopsis, 329
Microlepia, 568, 585; *brasiliensis*, 585; *cristata*, 585; *flaccida*, 586; *hirta*, 586; *japonica*, 586; *Khasiyana*, 586; *Mannii*, 582; *polypodioides*, 586; *Speluncae*, 585, 586; *strigosa*, 585; *tenuifolia*, 585
Mitosis in living cells, 407
Monascus, 149, 150
Monolepis spathulata, 311
Montia Viae, 314
Mucor Mucedo, 456, 457, 464-466, 471, 473-478, 482, 484-486, 488, 489, 492-495, 497; *stolonifera*, 456, 457, 464-469, 471-478, 482-489, 492-497
Mucor, protoplasmic streaming in, 455
Muhlenbergia, 102; *curtifolia*, 102; *neomexicana*, 102; *pauciflora*, 102; *polycaulis*, 102; *Pringlei*, 102
Muscites Stolzii, 342
Myrica emarginata, 390, 393; *longa*, 388
Myricales, 393
Myrtales, 402

Nantucket, The ferns and flowering plants of, 69, 415
Nasturtium trachycarpum, 322
Neottopteris, 569
Nephrolepis, 568, 583; *cordifolia*, 583, 584; *cultrifolia*, 584; *exaltata*, 584; *pendula*, 583; *tuberosa*, 583, 584
Nephrodium cyatheoides, 599; *Dubruleilianum*, 599; *Ecklonii*, 598; *exaltatum*, 584; *globuliferum*, 591; *Hudsonianum*, 601; *Keraudrenianum*, 596; *latifrons*, 593; *molle*, 599; *nymphale*, 599; *parasiticum*, 599; *polycarpon*, 600; *propinquum*, 598; *punctatum*, 578; *resiniferum*, 598; *rubiginosum*, 594; *squamigerum*, 594; *truncatum*, 601; *unidentatum*, 597; *violascens*, 599
Neurogramme calomelanos, 588
New ferns from Tropical America, 285
New West Indian *Lejeuneae*—II, 603
Nitella, 467
Notes on the genus *Widdringtonites*, 341
Notes on two Jamaica plants, 12
Notes on the North American species of *Phanerophlebia*, 23
Notes on species of *Solanum*, 11
Nothochlaena ternifolia, 576

Obione phyllostegia, 312; *tetraptera*, 311
Observations on the degree of stomatal movement in certain plants, 15
Ocotea, 401
Octoclinis, 342
Odontoloma, 568, 582; *Macraeanum*, 582, 583; *pulchellum*, 582
Odontosoria, 568, 584; *chinensis*, 585
Oenothera, 19, 22, 519-521, 527, 531; *biennis*, 15, 21, 520, 527, 528, 532; *grandiflora*, 519-527, 532; *Lamarckiana*, 519-532; *suaveolens*, 520, 522, 524, 527-529, 531
Olfersia aemula, 571; *gorgonea*, 572
Onagra vulgaris, 527
Onoclea, 259, 273; *Boryana*, 378; *lineata*, 366; *polypodioides*, 271, 273; *sorbifolia*, 259; *Spicant*, 275; *striata*, 375
Ophioderma, 234, 235; *pendulum*, 235
Ophioglossaceae, 234
Ophioglossales, 233
Ophioglossum, 234; *concinnum*, 234, 235; *ellipticum*, 235; *nudicaule*, 235; *pendulum*, 235; *vulgatum*, 234
Opuntia convexa, 290; *Griffithsiana*, 291; *leptocarpa*, 289; *Lindheimeri*, 289, 290; *macrorrhiza*, 289, 290; *reflexa*, 292; *texana*, 290
Opuntia Lindheimeri, Three new species of *Opuntia*, with a discussion of the identity of, 289
Oreodaphne, 400; *alabamensis*, 390, 400, 406
Ornithopteris aquilina, 575
Oryzopsis cuspidata, 102
Osmunda, 258, 259; *borealis*, 275; *cinnamomea*, 258; *Claytoniana*, 258; *lineata*, 366; *Lunaria*, 234; *polypodioides*, 271; *procera*, 383; *regalis*, 258; *Spicant*, 257, 258, 273, 275; *Struthiopteris*, 258, 259
Oxalidaceae, 417
Oxalis cymosa, 417; *stricta*, 417, 418
Oxycoccus, 551

Paesia aquilina, 575
Palaeocassia laurinea, 390, 396
Paliurus membranaceus, 401
Panicum nitidum, 102
Papilionaceae, 69
Parietaria occidentalis, 106; *pennsylvanica*, 106
Parrya macrocarpa, 326; *platycarpa*, 326
Parthenocissus, 39, 50

- Pellaea, 567, 575, 579; ternifolia, 576;
 Weddelliana, 576
 Phaeopeziza Nuttallii, 171
 Phanerophlebia, 23; auriculata, 24, 26;
 guatemalensis, 23, 26, 28; juglandi-
 folia, 23-25; pumila, 23, 25; macrosora,
 24, 26; nobilis, 24, 25; remotispora, 23,
 25; umbonata, 23, 26
 Phanerophlebia, Notes on the North
 American species of, 23
 Phegopteris crinalis, 595; Hillebrandii,
 595; honolulensis, 595; Keraudreniana,
 596; Keraudreniana procera, 596;
 microdendron, 600; punctata, 578;
 punctata flaccida, 579; punctata glabra,
 578; sandwicensis, 598; spinulosa, 597;
 unidentata, 597
 Phlebodium, 569
 Phragmidium, 120; Potentillae-canaden-
 sis, 114, 122; speciosum, 113, 115, 120,
 122; violaceum, 113, 119, 127, 128
 Phycomyces nitens, 456, 457, 464, 466,
 471, 488, 492, 494
 Phyllites aristolochiaeformis, 388; rhom-
 boideus, 388
 Phymatodes, 569
 Physaria didymocarpa lanata, 322;
 lanata, 322
 Physcia parietina, 201
 Picea alba, 99; albertiana, 99; canadensis,
 99
 PICKETT, F. L. A case of changed
 polarity in Spirogyra elongata, 509
 Picrococcus, 549, 553; elevatus, 559;
 floridanus, 554, 556
 Pilobolus, 170, 186
 Pinaceae, 99
 Pinonia, 242; splendens, 243
 Pinus Mertensiana, 100
 Plagiogyra, 257
 Platanus primaeva, 388
 Plicaria violacea, 170
 Poa, 104; alpicola, 104; ampla, 104;
 andina, 105; arctica, 105; arida, 104,
 105; brevipaniculata, 105; Buckleyana,
 104, 105; caesia strictior, 105; cali-
 fornica, 105; callichroa, 105; crocata,
 104, 105; Fendleriana, 105; flava, 104;
 flexuosa occidentalis, 105; Grayana,
 104; idahoensis, 105; interior, 105;
 juncifolia, 105; Kingii, 106; laevigata,
 105; laxa, 104; longiligula, 105; Mult-
 nomae, 104; nemoralis, 105; nervosa,
 105; nevadensis, 105; occidentalis, 105;
 paddensis, 104; Pattersonii, 104; phoe-
 nicea, 104; pratensis, 104; pratericola,
 105; pseudopratensis, 104; purpuras-
 cens, 104; rupicola, 105; Sandbergii, 104;
 serotina, 104; Sheldonii, 105; subpur-
 purea, 104; tenuifolia, 104; Tracyi, 105;
 triflora, 104
 Poaceae, 102
 Podophyllum, 441
 Podozamites lanceolatus, 390, 391
 Polyalthia, 502, 505
 Polycodium, 549, 551-553, 556, 557,
 caesium, 555, 556; candicans, 555, 558;
 elevatum, 555, 558; floridanum, 555;
 556, 558; **Kunthianum**, 559; Langloisii,
 555, 558; melanocarpum, 555, 557, 558;
 neglectum, 555, 557; oblongum, 555-
 558; oliganthum, 555; revolutum, 555,
 556; **sericeum**, 559; stamineum, 552,
 555, 557, 558
 Polygala cruciata, 421; polygama, 421;
 verticillata, 421; viridescens, 421
 Polygalaceae, 421
 Polygonaceae, 306
 Polypodiaceae, 237
 Polypodium, 285, 568; cordifolium, 583;
 crinale, 595; dentatum, 599; dictyo-
 tomum, 240; exaltatum, 584; Filix-mas,
 589; glaucum, 239; Hillebrandii, 595;
 honolulense, 595; **insidiosum**, 287, 288;
 Keraudrenianum, 596; lineare, 240;
 molle, 599; nymphale, 599; Paepigii,
 578; parasiticum, 598; pectinatum, 239;
 polycarpon, 600; procerum, 596; punc-
 tatum, 578; sandwicense, 598, 600;
 Speluncae, 586; spinulosum, 597; steg-
 nogrammoides, 600; trifurcatum, 285,
 287; unidentatum, 597
 Polystichum, 569; Dubrueilianum, 599;
 molle, 599; truncatum, 601
 Populus, 301, 388; acuminata, 302;
 balsamifera, 301; balsamifera candi-
 cans, 302; Besseyana, 302; candicans,
 301; Fremontii, 303; harkeriana, 390,
 394; hastata, 301; Sargentii, 301, 302;
 trichocarpa, 301
 Porella, 280, 329; platyphylla, 280
 Portlandia **Harrisii**, 8; **nitens**, 10
 Portulacaceae, 314
 Potamogeton perfoliatus, 101; Richard-
 sonianus, 101
 Prinos padifolius, 426
 Proteoides daphnogenoides, 394
 Protoplasmic streaming in Mucor, 455
 Pseudocarpidium avicennioides, 10; **pung-**
ens, 10; **rigens**, 10
 Pteridophyta, 232
 Pteridophyta of the Hawaiian Islands.
 A taxonomic study of the, 227, 567
 Pteridium, 567, 574; aquilinum, 575
 Pteris, 567, 573, 579; alata, 574;
 aquilina, 575; arachnoidca, 575;
 arborea, 573; atropurpurea, 576; bi-
 aurita, 579; cretica, 573; decipiens,
 576; decora, 577; excelsa, 573; irregu-
 laris, 573, 574; lomarioides, 574;
 Lydgatei, 579; melanocaulon, 574;
 nervosa, 573; owahuensis, 573; pedata,
 576; pentaphylla, 574; peruviana, 576;
 psittacina, 575; regularis, 574; scab-

- ripes, 574; serraria, 573; subverticillata, 576; terminalis, 573; ternifolia, 576; Treacheriana, 574; triphylla, 574
- Puccinia Bunii, 117; Cirsii-lanceolati, 114-116, 124; Liliacearum, 117; Malvacearum, 128, 129; Peckiana, 116; Podophylli, 114; transformans, 114, 115
- Pyronema, 63-66, 146, 147, 149, 150, 190, 191; omphalodes, 67
- Pyronema, The viability of the spores of, 63
- Radicula *trachycarpa*, 322
- Radula, 210, 329, 338; complanata, 329, 331, 332, 336, 338; flaccida, 329-339; Hedingeri, 329, 334, 335, 337, 338, 339; Lindbergiana, 329, 338; protensa, 329, 336, 339; subtropica, 329, 338; tjibodensis, 329, 338
- Radula, Discoid gemmae in, 329
- Ramalina farinacea, 201; fraxinea, 201; scopulorum, 201
- Ranales, 395
- Ranunculaceae, 319
- Ranunculus Macounii, 319; repens, 319; *rivularis*, 319
- Rectolejeunea, 611; Brittoniae, 610, 611; flagelliformis, 279; *Maxonii*, 609, 611; phyllobola, 611
- Relation of Ohio bog vegetation to the chemical nature of peat soils, The, 53
- Rhamnales, 398
- Rhamnus tenax, 390, 398
- Rhizopus nigricans, 457
- Rhus, 398; copallina, 423; cretacea, 397; glabra, 423; hirta, 423; reddita, 397; *redditiformis*, 390, 397, 406; Uddeni, 397; villosa, 398
- Rhyarobius, 142, 145, 148, 149, 186, 193; albidus 143, 148; Cookei, 143, 148; crustaceus, 152; niveus, 152; pachyascus, 143, 148, 152; sexdecimsporus, 152
- Robinia Pseudacacia, 74; viscosa, 74
- ROBINSON, C. B. Polycodium, 549
- ROBINSON, W. J. A taxonomic study of the Pteridophyta of the Hawaiian Islands, 227, 567
- Rondeletia *saxicola*, 8
- Rosales, 396
- Ruppia curvicaarpa, 101; maritima, 101
- RydbERG, P. A. Studies on the Rocky Mountain flora, 99, 301
- Sabina *horizontalis*, 100; prostrata, 100
- Saccobolus, 190; depauperatus, 152; Kerverni, 152; neglectus, 152, 168, 169, 193; violascens, 152
- Sadleria, 569
- SAFFORD, W. E. Desmos the proper generic name for the so-called Unonas of the Old World, 501
- Sagittaria, 101; arifolia, 102; arifolia tenuior, 102; cuneata, 102; paniculata, 101, 102
- Sagina occidentalis, 317
- Salicaceae, 301
- Salicales, 394
- Salix, 387, 388, 439; alexensis, 304; arbusculoides, 304; arguta erythrocoma, 304; Barrettiana, 304; bella, 304; conjuncta, 304; cordata, 304; cordata angustata, 303; *cordata* \times *monticola*, 303; deleta, 388; desertorum, 304; Drummondiana, 304; erythrocoma, 304; Fernaldii, 304; fragilis, 439; glaucops, 303; *glaucops* \times *monticola*, 303; Hookeriana, 304; lasiolepis, 304; lucida, 304; MacCalliana, 304; monticola, 303; petiolaris, 49; *Sandbergii*, 304; saskatchewanana, 304; Seemanii, 304; sitchensis, 304
- Salpichlaena, 257
- Salviniaceae, 232
- Salviniales, 232
- Sapindales, 396
- Sapindus Morrisoni, 388, 390, 396
- Sapranthus, 502; nicaraguensis, 502
- Schizaea, 237; australis, 238; dichotoma, 237; robusta, 238
- Schizostegia, 568, 579; Lydgatei, 579; calocarpa, 579; pachysora, 579
- Schizaeaceae, 237
- SCHREINER, O., and SKINNER, J. J. The effect of guanidin on plants, 535
- SEEVER, F. J. Studies in pyrophilous fungi—III. The viability of the spores of Pyronema, 63
- Sequoia, 389
- Seubertia, 111
- Sexual fusions and spore development of the flax rust, 113
- Silene, 317; Drummondii, 318
- Simarubaceae, 420
- Sisymbrium officinale, 323
- Sitanion, 106; brevifolium, 106; ciliatum, 107; elymoides, 107; insulare, 107; lanceolatum, 107; longifolium, 106; marginatum, 107
- SKINNER, J. J., and BEATTIE, J. H. Effect of asparagin on absorption and growth in wheat, 429
- SKINNER, J. J., and SCHREINER, O. The effect of guanidin on plants, 535
- SLOSSON, M. New ferns from Tropical America, 285
- Smelowskia americana, 327; *lobata*, 327; ovata, 327
- Solanum bahamense, 11; bahamense subarmatum, 11; Blodgettii, 11; boldoense, 11; verbascifolium, 11
- Solanum, Notes on species of, 11
- Sophia *leptostylis*, 325; procera, 325
- Sordaria, 190

- Sparganiaceae, 101
 Sparganium angustifolium, 101; longipedunculatum, 101; simplex, 101
 Sphaerotherca, 149
 Spicanta attenuata, 272; borealis, 275; Ghiesbreghtii, 277; L'Herminieri, 268; onocleoides, 271; polypodioides, 272
 Spirogyra communis, 509, 510; elongata, 509
 Spirogyra elongata, A case of changed polarity in, 509
 Sporobolus vaginaefolius, 103
 Stegania, 257, 259
 Stegogramma sandwicense, 600
 Stellaria alpestris, 315; borealis, 315; longipes, 315; stricta, 315; subvestita, 315; valida, 315
 Stemphylium, 142
 Stenanthella occidentalis, 108
 Stenochlaena, 257, 372
 Stenoloma tenuifolium, 585
 Sterculia lugubris, 390, 399, 406
 Stibasia Douglasii, 236
 Stictolejeunea, 211
 Stipa hymenoides, 102; membranacea, 102
 Stropholirion, 111; californicum, 111
 Struthiopteris, 257-259, 261, 264, 364, 373, 377; **chiriquana**, 359-362, 380; **Christii**, 360, 362, 380, 382; **costaricensis**, 360, 363, 364, 372, 384; **danaeacea**, 359, 364, 380; doodioides, 259, 275; ensiformis, 261, 262; **exaltata**, 262, 264, 266, 278; **falciformis**, 357, 365, 359, 365; **jamaicensis**, 262, 266; **L'Herminieri**, 262, 267; **lineata**, 360, 366, 368, 369, 374-377, 381-384; **Maxonii**, 262, 268; ornifolia, 373; pennsylvanica, 258; **Plumieri**, 261, 262, 266, 269, 270, 271; polypodioides, 259, 261, 270, 271, 275, 364, 383; **rufa**, 357, 358, 369; **Schiedeana**, 359, 370, 377, 380, 385; **sessilifolia**, 360, 362, 373; **Shaferi**, 357, 358, 374, 375, 378; **Spicant**, 258, 259, 262, 277, 278; **stolonifera**, 262, 277, 278; **striata**, 359, 360, 368, 369, 375-377, 380-385; **Underwoodiana**, 357-359, 375, 377, 380, 385; **varians**, 357, 359, 379, 380; **violacea**, 359, 373, 376, 378-382; **vivipara**, 359, 380-382; **Werckleana**, 357, 358, 364, 373, 382, 384
 Struthiopteris and its representatives in North America, The genus, 257, 357
 Studies in pyrophilous fungi—III. The viability of the spores of Pyronema, 63
 Studies on the Rocky Mountain flora, 99, 301
 Studies of West Indian plants—IV, 1
 Symplocarpus foetidus, 439
 Tabernaemontana discolor, 12; ochroleuca, 12
 Taxonomic study of the Pteridophyta of the Hawaiian Islands, A, 227, 567
 Tecoma stans, 115
 Teloschistes chrysophthalmus, 201; flavicans capensis, 201
 Tetraclinis, 342
 Texas. Contributions to the Mesozoic flora of the Atlantic coastal plain—VIII, 387
 Tectaria, 569
 Thalictrum **columbianum**, 320; megacarpum, 320; venulosum, 320
 Thecotheus, 187; Pelletieri, 146, 152, 164, 166, 193
 Thelebolus, 139, 186; stercoreus, 143, 146-149, 152, 167, 169, 170
 Thuia graminea, 342
 Thuites gramineus, 342
 Thymeleales, 399
 Thyopsiella, 280
 Tofieldia coccinea, 108; occidentalis, 108
 Toxicodendron Vernix, 424; vulgare, 424
 Toxicoscordion, 110
 Trachycolea, 280
 Tradescantia, 107; bracteata, 108; laramiense, 107; occidentalis, 107; scopulorum, 108; universitatis, 107; virginica occidentalis, 107
 Tricalycites papyraceus, 390, 405
 Trichomanes, 244; apiifolium, 245; Bauermanum, 244, 245; bifolium, 245; chinensis, 585; cuneiforme, 595; crispum, 244; cyrtotheca, 245, 246; davallioides, 246; Draytonianum, 246; eminens, 245; Endlicherianum, 246; humile, 245, 246; meifolium, 245; minutulum, 246; minutum, 245; myrioplazium, 245; parvulum, 244, 245; radicans, 245, 246; sandwicense, 246; saxifragoides, 245; sibthorpioides, 245; speciosum, 246; strigosum, 585
 Trifolium arvense, 73; aureum, 72; dubium, 73; hybridum, 73; incarnatum, 73; pratense, 73, 74; procumbens, 73; repens, 74
 Trillium, 439
 Triphragmium, 114-116; Ulmariae, 113, 114, 122
 Tritelia, 111
 Tsuga, 100; Mertensiana, 100; Pattoniana, 100
 Tumion, 389
 Ulex europaeus, 69
 Ulmaceae, 304
 Umbellales, 402
 Undescribed species from Cuba, 9
 Undescribed species from Jamaica, 7
 Unona, 501-504; acutiflora, 502; bibracteata, 501; Brandesana, 504; cochinchinensis, 506; crassipetala, 502; crinita, 505, 507; dasymaschala, 507;

- desmantha, 505, 508; Desmos, 504, 506; discolor, 502, 504, 506; discreta, 502, 503; dumosa, 506; Dunalii, 506; elegans, 506; latifolia, 504; Lawii, 506; longiflora, 507; nitidissima, 502; panamensis, 501; pannosa, 506; penduliflora, 502; praecox, 507; stenopetala, 505, 507; uncinata, 502; violacea, 502; viridiflora, 506; Wrayi, 505, 507; xylopioides, 502; zeylanica, 506
- Unonopsis, 501
- Uromyces Caladii, 113; Lillii, 114; Scirpi, 114, 115
- Urtica Breweri, 305; cardiophylla, 306; gracilis, 305, 306; **strigosissima**, 305; **viridis**, 305
- Urticaceae, 305
- Urticales, 394
- Usnea articulata, 201; floridana, 201
- Uvaria latifolia, 504; monilifera, 502
- Uvularia sessilifolia, 439-441
- Uvularia sessilifolia, A contribution to the life history of, 439
- Vaccinium, 549-554, 556; album, 553; arboreum, 549, 551, 553; caespitosum, 552, 556; confertum, 551; consanguineum, 554, 555; corymbosum, 552; crassifolium, 553; cubense, 552; dumosum, 553; elevatum, 553, 559; frondosum, 553; Kunthianum, 552, 553, 555, 559; leucanthum, 552; melanocarpum, 559; melanocarpum candicans, 557; melanocarpum sericeum, 557-559; meridionale, 554; multiflorum, 554; myrtifolium, 553; Myrtillus, 552, 556; pallidum, 553; poasanum, 551; Ramonii, 554; scoparium, 556; stamineum, 549, 553-555, 559; stamineum melanocarpum, 557; stenophyllum, 552; Vitis-Idaea, 551, 553
- Variation in the floral structures of Vitis, 37
- Vatica, 406
- Vegetative reproduction in the New England Frullaniaceae, 279
- Veratrum Eschscholtzianum, 108
- Viburnum robustum, 388, 390, 405
- Vicia angustifolia, 79; hirsuta, 78; sativa, 79; tetrasperma, 78; villosa, 78
- Viola, 92, 93; Angellae, 88-90; fimbriatula, 92; hirsutula, 95; **hirsutula** × **palmata**, 96; hirsutula × sororia, 96; **hirsutula** × **Stoneana**, 96; **hirsutula** × **triloba**, 95; latiuscula, 95; **latiuscula** × **triloba**, 94, 95; palmata, 85-89, 97; palmata dilatata, 92, 96; palmata × papilionacea, 85, 97; **palmata** × **triloba**, 88-90, 96; palmata × villosa, 95; papilionacea, 85-93, 97; **papilionacea** × **Stoneana**, 93; **papilionacea** × **triloba**, 85, 90-92; pedatifida, 97; sagittata, 92; sororia, 90, 92, 93; **sororia** × **triloba**, 92; Stoneana, 93; **Stoneana** × **triloba**, 93, 96; Stoneana × villosa, 96; triloba, 88-97; variabilis, 89, 90, 91; villosa × palmata asarifolia, 96
- Violet hybrids between species of the palmata group, 85
- Viscaria, 317
- Vitex rigens, 10
- Vitis, 37-39, 46, 50, 51; aestivalis, 43; bicolor, 38, 40, 43, 48, 49; Labrusca, 40, 48; vinifera, 39, 42, 48; vulpina, 38-45, 48, 49, 51
- Vitis, Variation in the floral structures of, 37
- Vitis, General floral habits of, 37; Special variation in flower parts, 42
- Vittaria, 568, 586; elongata, 587; rigida, 587
- Wahlbergella, 317, 318; affinis, 318; apetala, 319; **attenuata**, 319; **Drummondii**, 318; **Kingii**, 319; **montana**, 318; **Parryi**, 319; **striata**, 318; **Taylorae**, 318; triflora, 318; uniflora, 319
- Was Lamarck's evening primrose (Oenothera Lamarckiana Seringe) a form of Oenothera grandiflora Solander?, 519
- West Indian Lejeuneae, New—II, 603
- West Indian plants, Studies of —IV, 1
- West Indies, Dendropanax in the, 1
- Widdringtonia, 342, 343
- Widdringtonites, 342-344, 346, 347; oblongifolius, 346; Reichii, 346; subtilis, 344, 346, 347; Ungeri, 342
- Widdringtonites, Notes on the genus, 341
- Williston, R. Discoid gemmae in Radula, 329
- Woodwardia, 276
- Xylopia, 504; frutescens, 504; grandiflora, 502; muricata, 504; salicifolia, 504
- Zamia lanceolata, 391
- Zannichelliaceae, 101
- Zizyphus, 398; **lamarensis**, 390, 398, 406
- Zygadenus alpinus, 108; dilatatus, 108; elegans, 108; falcatus, 110; glaberrimus, 110; gramineus, 109; intermedius, 109, 110; Nuttallii, 110; paniculatus, 110; porrifolius, 108; venenosus, 109
- Zygnema pectinatum, 128